



Animal Clinic, LLC

232 First Street
Fort Lupton, CO 80621
303 857-6671
303 857-6673 FAX

BEEF COW VACCINATION PROTOCOL FOR MARCH CALVING HERD

Baby Calves at birth: (March)

- Colostrum – within 6 hours of birth.
- Dip navels in strong Iodine (7%).
- Band bulls not intended for breeding
- C & D Antitoxin
- *Bulls that were banded should also receive a tetanus vaccine

Baby Calves at branding time: (1-3 months of age): (April - May)

- Brand
- Dehorn
- Modified live respiratory – Bovishield Gold
- 7 or 8 way Clostridium – Ultrabac
- *Castrate bulls that were not banded and not intended for breeding

Baby Calves - Booster dose of vaccine (4 weeks after branding vaccines): (May - June)

- Modified live respiratory – Bovishield Gold
- 7 or 8 way Clostridium – Ultrabac

Two weeks prior to weaning: (October - November)

- Modified live respiratory – Bovishield Gold OneShot
- 7 or 8 way Clostridium – Ultrabac
- Bangs vaccinate heifers – Veterinarian
- Parasite Control – An Avermectin pour on

Fall pregnancy check: (October – November)

- Cows and Bulls:
 - Pregnancy check cows
 - Parasite Control – An Avermectin pour on
 - 7 or 8 way Clostridium – Ultrabac
 - Calf Scours vaccine – Scourguard 3k

Late Fall Booster dose of Scours vaccine (3-4 weeks after initial dose and at least 6 weeks before calving):

- Calf Scours vaccine – Scourguard 3KC

Spring – after calving and before breeding: (April - May)

- Cows and Bulls:
 - Modified live respiratory – Bovishield Gold
 - Vibrio Lepto 5 vaccine – Tri-Vib 5L
- Yearling replacement heifers:
 - Modified live respiratory – Bovishield Gold
 - Vibrio Lepto 5 vaccine – Tri-Vib 5L
 - 7 or 8 way Clostridium – Ultrabac