



Animal Clinic, LLC 232 First Street Fort Lupton, CO 80621 303-857-6671

Dr. Ray Sagehorn, Dr. Troy Dowdy, Dr. Victoria J. Cabrera, Dr. Samantha Delbecq, Dr. Sawyer Howell

EQUINE VACCINATION GUIDELINES: BROODMARES AND STALLIONS

Vaccine	Broodmares	Stallions	Comments
Tetanus Toxoid*	Annual Give 4-6 weeks prior to foaling	Annual Give prior to breeding season	Re-vaccinate after penetrating wound or surgery if last dose was greater than 6 months
Eastern & Western * Encephalomyelitis	Annual Give 4-6 weeks prior to foaling	Annual Give prior to breeding season	
Rabies*	Annual Give 4-6 weeks prior to foaling	Annual Give prior to breeding season	State regulations require this vaccine to be given by a licensed veterinarian for the horse to be classified as vaccinated
West Nile Virus*	Annual Give 4-6 weeks prior to foaling	Annual Give prior to breeding season	Consider booster vaccine in Fall if mosquito load is significant
Equine Herpes Virus 1 & 4	Give at 5, 7, and 9 months of gestation. Include in the 4-6 weeks prior to foaling	Semi-annual Give prior to and after breeding season	
Equine Influenza Virus	Semi-annual Give 4-6 weeks prior to foaling	Semi-annual Give prior to and after breeding season	
Strangles	*Vaccinate 4-6 weeks prior to foaling with approved killed products	*Vaccinate every 6-12 months	*Vaccination should be based on risk-assessment and consulted with your veterinarian
Rotavirus	Give at 8, 9, and 10 months of gestation	Not applicable	

*These are considered CORE vaccines by the American Association of Equine Practitioners and should be apart of every horses yearly vaccine rotation.

Regional variations in vaccines recommendations exists. Depending on the region of the country, some vaccines not listed here may be necessary. Also the frequency of these vaccines may vary depending on exposure in certain instances. Consult with your veterinarian regarding these variation potentials.